

Note to Readers: If you need assistance accessing items in this Supplemental Material, please contact ehp508@niehs.nih.gov. Our staff will work with you to assess and meet your accessibility needs within 3 working days.

Table of Contents for Supplemental Material

Anthropometry in 5- to 9-Year-Old Greenlandic and Ukrainian Children in Relation to Prenatal Exposure to Perfluorinated Alkyl Substances

Birgit Bjerre Høyer, Cecilia Høst Ramlau-Hansen, Martine Vrijheid, Damaskini Valvi, Henning Sloth Pedersen, Valentyna Zvezdai, Bo A.G. Jönsson, Christian H. Lindh, Jens Peter Bonde, and Gunnar Toft

Table S1. Characteristics of participants and those lost to follow-up.

Table S2. Maternal PFOA and PFOS concentrations during pregnancy and crude relative risk (RR) of offspring overweight (WHO > 85th percentile (age and sex standardised)) and waist-to-height-ratio >0.5 at 5 to 9 years.

Table S3. Maternal PFOA and PFOS concentrations during pregnancy and crude and adjusted associations with BMI z-scores (sex and age standardised) and WHtR in offspring at 5 to 9 years.

Table S4. Complete-case-analysis of maternal PFOA and PFOS concentrations during pregnancy and crude and adjusted relative risk (RR) of offspring overweight (WHO > 85th percentile (sex and age standardised)) and waist-height-ratio >0.5 at 5 to 9 years.